

Reglan Side Effects

The most extreme side effects of Reglan generally stem from prolonged use or higher doses, and are described by victims as embarrassing and debilitating. Reglan may worsen symptoms of Parkinson's disease, depression or bowel obstructions.

Reglan side effects include:

- Drowsiness
- Shuffling walk
- Constipation
- Dizziness
- Drooling
- Depression
- Tiredness
- Tremors
- Tardive dyskinesia (see below)
- Trouble swallowing
- Headache
- Fever
- Stiff muscles
- Confusion
- Sweating
- Severe anxiety
- Irregular heartbeat
- Rapid breathing
- Suicidal thoughts
- Hallucinations
- Mental/mood changes
- Seizure
- Jaundice

Tardive Dyskinesia

Patients are most prone to tardive dyskinesia, or uncontrolled muscle movements in the face, tongue and extremities, when Reglan is taken for more than 12 weeks or if they are elderly.

Reglan Side Effects (Cont.)

Tardive dyskinesia cannot be treated, but occasionally the cessation of the medicine intake can decrease or eliminate the symptoms.

Reglan Lawyers

If you or someone you love has suffered serious harm as a result of taking Reglan, you may be eligible to seek compensation, including medical bills, lost earning capacity, pain and suffering, and more. Contact the Simmons firm today to get in touch with caring and experienced Reglan lawyers who can evaluate your case, answer your questions, and help you understand your legal rights, including the filing of a Reglan lawsuit.